

then measured outside the tank. The results are used to calculate body volume and body density. The body density is then used to calculate the proportion of body weight as fat mass (FM) and fat free mass (FFM).^[29] Recent developments of this method use air rather than water displacement, in a device called a BOD-POD. An

exciting development in the application of air-displacement plethysmography in the testing of infants (< 1 year old) has recently emerged. Life Measurement Incorporated has developed an air-displacement plethysmograph specifically for infants between birth and 6 months of age with body masses of 2.7-7.4 kg.^[30]

Table 1. Prevalence of pediatric obesity worldwide

Country	Date of survey	Age (years)	Prevalence (%)	References
South America				
Argentina	2000	2-5	7.2	de Onis & Blossner, 2000 ^[8]
Bolivia	1998	2-5	6.5	de Onis & Blossner, 2000 ^[8]
Brazil	1996 1997	2-5 6-18	8 14 (overweight + obesity)	de Onis & Blossner, 2000 ^[8] Wang et al, 2002 ^[6]
Chile	1996 2000	0-5 6-19	7 7 (m), 8 (f)	de Onis & Blossner, 2000 ^[8] Kain et al, 2002 ^[7]
Columbia	1995	2-5	2.5	de Onis & Blossner, 2000 ^[8]
Mexico	2000 2003	0-5 <5	4.9 5.5	Gonzales et al, 2004 ^[9] Avila et al, 2003 ^[10]
Peru	1996	2-5	6	de Onis & Blossner, 2000 ^[8]
Uruguay	2000	2-5	6.1	de Onis & Blossner, 2000 ^[8]
Venezuela	1997	2-5	3	de Onis & Blossner, 2000 ^[8]
North America				
Canada	1996	7-13	10 (m), 9 (f)	Tremblay et al, 2002 ^[11]
USA	1988-1994	6-11	11	Troiano et al, 1995 ^[12]
	1988-1994	12-17	13 (m), 9 (f)	Flegal et al, 2001 ^[18]
		6-11 12-17	8 (m), 7 (f) 8	
Europe				
Austria	2002-2003	10-15	9 (m), 5 (f)	Widhalm and Dietrich (personal communication, 2003)
France	1990	4-17	3	Rolland-Cachera et al, 2002 ^[14]
Germany	1995	7-14	8 (m), 10 (f)	Kromeyer-Hauschild et al, 1999 ^[15]
Greece	1998 2000	12 6-17	18 (m), 5 (f) 4	Mamalakis et al, 2000 ^[16] Krassas et al, 2001 ^[17]
Italy	1994-2000	6-20	4	Cacciari et al, 2002 ^[18]
Portugal	2003	7-10	9 (m), 12 (f)	Padez et al, 2003 ^[19]
Russia	1998	6-18	9	Wang et al, 2002 ^[6]
Spain	1995 1995	6-15 6-15	6 (m), 2 (f) 7	Moreno et al, 2000 ^[20] Rios et al, 1999 ^[21]
Sweden	1997	12-18	8 (m), 4 (f)	Berg et al, 2001 ^[22]
Switzerland	1999	6-12	10	Zimmermann et al, 2000 ^[23]
UK	1998	7-11	5 (m), 4 (f)	Lobstein et al, 2003 ^[24]
Africa				
Algeria	1996	2-5	9	de Onis & Blossner, 2000 ^[8]
Egypt	1996 1997	2-5 10-19	8.5 4	de Onis & Blossner, 2000 ^[8] Ibrahim et al, 2002 ^[25]
Iran	1995	2-11	3	de Onis & Blossner, 2000 ^[8]
Marocco	1992	2-5	6.5	de Onis & Blossner, 2000 ^[8]
S. Africa	1996	2-5	6.5	de Onis & Blossner, 2000 ^[8]
Tunisia	1988	2-5	4	de Onis & Blossner, 2000 ^[8]
Asia-Pacific region				
Australia	1995	2-18	5	Magarey et al, 2001 ^[26]
China	1995	6-18	8 (m), 7 (f)	Ke-You & Da-Wei, 2001 ^[27]
India	2002	13-18	4 (m), 3 (f)	Ramachandran et al, 2002 ^[28]

Countries selected with all information available.

Bioimpedance analysis

Bioimpedance analysis (BIA) is based on measurement of electrical resistance in the body to a tiny imperceptible current. This approach provides an estimate of total body water, which can be transformed into fat free mass. The main concept of BIA is that tissues rich in water and electrolytes are less resistant to the passage of an electrical current than lipid-rich adipose tissue. BIA methods require use of appropriate equations (age and population specific), standard measurement conditions, and several other factors (body position, time of the day, and room temperature).^[27,31] Prediction equations should only be applied to the population for whom the regression equation has been developed.

Dual energy X-ray absorptiometry

Dual energy X-ray absorptiometry (DXA) quantifies the relative attenuation of two main photon peaks as they pass through tissue. The relative attenuation of the two photon streams is a function of tissue elemental composition. This technique is more feasible with children for whom other laboratory techniques may be impractical (underwater weighing). The greatest advantage of DXA may be the ability to assess regional body composition (trunk, arms and legs). An example of a research use of DXA that may lead to clinical application is the prediction of co-morbidity risk in obese children and adolescents.^[32]

Imaging

Computerized axial tomography (CT) and magnetic resonance imaging (MRI) provide investigators with the opportunity to evaluate tissue-system level components *in vivo*.^[33] In particular, the effect of fat distribution on disease risk is a subject of great interest. In children and adolescents, recent studies have related central fat to Type 2 diabetes mellitus and cardiovascular disease.^[34] Imaging can inform the origins of the relationships between visceral adipose tissue and adverse health and also explain the influence of ethnic and gender differences in adipose tissue distribution.^[35]

Clinical setting methods

Anthropometry is the least expensive and most widely used method for assessing pediatric body composition, and is especially informative for epidemiological studies.

Body mass index

Body mass index (BMI) is defined as weight (kg)/height squared (m^2) and is widely used as an index of relative adiposity. For children, various cut-off criteria have been proposed based on reference populations and statistical approaches. BMI varies with age and gen-

der. It typically rises during the first month after birth, falls after the first year and rises again around the six years of life (adiposity rebound). A given value of BMI therefore needs to be evaluated against age- and gender-specific reference values. Children with a BMI greater or equal to the 95th percentile for age and gender are generally classified as "overweight" and children with a BMI between the 85th and 95th percentile as "at risk of overweight".^[36] Using data from different nations, Cole et al.^[37] derived curves that passed through the points of 25 and 30 BMI at age 18 years. These were used to provide age and gender specific BMI cut offs to define childhood overweight and obesity.

Skinfolds

Skinfold thickness measurements are widely applied in pediatric subjects, using special calipers to grasp a skinfold at different sites of the body (triceps, subscapular). Prediction equations can be used to estimate fat mass and fat percentage from the skinfold measurements.^[35]

Waist circumference

Waist circumference is an indirect measure of central adiposity, which itself is correlated with risk of cardiac disease and adverse lipid profile.^[38] Waist circumference is measured at the minimum circumference between the iliac crest and the rib cage using an anthropometric tape.^[39] Recently, normative percentiles have been published in the United States for waist circumference measures for children age 2 to 18 years in three major ethnic groups (African-American, European American, Mexican-American), using data from the Third National Health and Nutrition Examination Survey (NHANES III) (Table 2). These can be used to assess trends, compare individual children to population norms, and to compare populations to public health recommendations.^[40]

Comorbidities of childhood obesity

Childhood obesity is linked to multiple health risks including elevated blood pressure, respiratory abnormalities, dyslipidemia, hyperinsulinemia and health complications that are precursors to Type 2 diabetes.^[41]

Blood pressure

There is an association between adiposity and elevated blood pressure in children and adolescents.^[42-45] Elevated blood pressure occurs approximately nine times more frequently among obese than nonobese children.^[44] One

Table 2. Waist circumference according to sex for African-American, European-American and Mexican-American children^[40]

Age	European Americans				African Americans				Mexican Americans			
	Boys		Girls		Boys		Girls		Boys		Girls	
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
2	47.973	0.246	48.084	0.274	47.191	0.228	46.514	0.301	48.051	0.233	48.353	0.288
3	49.565	0.268	49.634	0.343	49.296	0.273	48.975	0.306	51.118	0.344	51.167	0.416
4	51.462	0.280	51.957	0.403	50.847	0.301	50.817	0.388	53.576	0.451	52.836	0.479
5	52.989	0.378	53.592	0.476	52.869	0.408	52.510	0.342	54.024	0.422	54.653	0.542
6	57.246	1.018	54.055	0.941	53.383	0.595	55.845	0.951	56.354	0.749	56.293	1.052
7	58.605	1.112	57.875	1.322	56.311	0.726	59.113	1.194	60.901	1.623	57.769	0.782
8	58.815	0.838	58.437	0.878	58.359	0.862	61.174	1.146	62.917	1.068	62.664	1.109
9	63.980	1.329	63.571	1.672	60.930	0.929	62.423	1.088	64.633	1.256	64.719	1.618
10	66.343	1.377	64.929	1.798	63.740	1.047	65.814	1.314	67.480	1.219	64.370	0.886
11	67.359	1.411	67.945	1.189	68.594	1.440	68.628	1.241	74.093	1.785	73.412	2.182
12	72.410	1.333	68.822	1.662	71.300	1.952	70.606	1.234	73.214	1.641	73.538	1.339
13	73.223	1.659	74.348	1.696	71.400	2.111	74.819	1.422	77.332	1.595	73.700	1.484
14	80.041	3.631	74.591	1.377	73.823	1.730	77.328	2.145	74.940	1.574	75.203	1.148
15	81.187	2.443	74.067	1.318	74.947	1.221	79.535	3.366	79.584	2.056	76.861	1.439
16	78.990	1.529	76.652	1.909	74.894	1.157	78.052	1.457	80.686	1.219	76.043	1.408
17	81.867	1.620	77.300	1.385	76.126	1.327	79.357	1.720	81.903	1.532	82.342	1.877
18	81.066	1.824	76.757	1.459	79.267	1.625	83.071	2.425	84.447	1.664	81.266	1.921

study examined data from over 3000 5 to 18 year old males and females^[45] and found that fatness in excess of 25% in males and 30% in females was associated with increased risk for hypertension. Similar findings have also been noted among 5- and 6-year-old Hispanic children.^[46] Daniels et al^[47] examined the predictors of left atrial enlargement in children and adolescents with essential hypertension, and found that BMI was significantly elevated among participants with vs without left atrial enlargement.

Respiratory abnormalities

In nonobese individuals, there is usually little fat deposition in the neck and in the peripharynx regions. Increased neck adiposity is associated with respiratory abnormalities during sleep among pediatric individuals. Respiratory complications associated with childhood obesity include asthma, obstructive sleep apnea, and pickwickian syndrome.^[48-49] Sleep apnea is especially common in childhood obesity.

Lipids and lipoproteins

Pediatric obesity is associated with elevated levels of total cholesterol, low-density lipoprotein (LDL) cholesterol, and triglycerides, and decreased levels of high-density lipoprotein (HDL) cholesterol.^[50-53] Freedman et al^[54] studied 10 to 17-year-old children from the Bogalusa Heart Study and found that children who were at risk for overweight or who were overweight were 2.4 times as likely as normal weight children to have elevated total cholesterol, diastolic blood

pressure, low-density lipoprotein cholesterol, high-density lipoprotein cholesterol, systolic blood pressure, triglycerides, and fasting insulin. Several of these associations differed between whites and blacks, and by age.

Glucose intolerance, insulin resistance, hyperinsulinemia, and Type 2 diabetes

Body fatness is a reliable correlate of hyperinsulinemia and glucose intolerance among adolescent males and females.^[55-59] It has been proposed that hyperinsulinemia may underlie the associations among obesity, hypertension, and glucose intolerance.^[60] These associations have been documented in pediatric samples.^[58,61] Impaired glucose tolerance is a condition indicated by elevated blood sugar levels, but not the levels high enough for a diagnosis of Type 2 diabetes. A clinic-based study found that 25% of obese children 4 to 11 years old, and 21% of obese children 11 to 18 years old, had impaired glucose tolerance.^[62]

Insulin resistance, another precursor to Type 2 diabetes, is also related to childhood obesity.^[63] In a school-based study of over 12 000 5th to 8th grade boys and girls, Sinaiko et al^[64] found significant associations between insulin resistance and body fat. The greatest insulin resistance was also associated with elevated triglycerides and LDL cholesterol, reduced HDL cholesterol, and elevated blood pressure.

The prevalence of Type 2 diabetes in children appears to be increasing. In an analysis of 1027 children and adolescents attending a Cincinnati clinic, only 4% were diagnosed with Type 2 diabetes mellitus before

1992.^[65] By 1994, 16% of new cases of Type 2 diabetes were diagnosed in that age group, with 10 to 19 years old representing 33% of all those cases. This translated into a 10-fold increase in the incidence of diabetes within the 10 to 19-year-old age group.

Treatment of child obesity

Family-based behavioral treatment programs have been the most extensively studied interventions to date and generally achieve the best short- and long-term results. Treatment strives to teach the family members the behavioral skills necessary to establish and sustain healthier eating and physical activity patterns. Behavior modification strategies such as behavioral contracting, stimulus control, and positive reinforcement therefore set the foundation to help children lose weight.^[66-68] Behavioral contracting refers to an explicit "contract" among family members that stipulates the behavioral goals that family members will attempt to reach and the reinforcements they will receive for attaining such goals. Rewards other than food and money are utilized. Family-based and interpersonal rewards are used instead (praise, family trips, sports equipment). Stimulus control refers to practical restructuring of the physical home environment such that healthier foods become more readily accessible whereas high-fat, high-sugar foods are less accessible. Positive reinforcement strategies for families form the foundation for treatment, as parents are trained to move away from punitive parenting strategies to more positive parenting strategies.

Research suggests that dietary modification is a powerful and necessary component for child weight loss. Treatments focusing solely on dietary modification have achieved short-^[69-70] and long-term weight losses^[71]. Short-term interventions lacking a dietary component achieved mixed results,^[72,73] but there is no evidence for long-term efficacy of treatments lacking a dietary component.

Many programs have used Epstein's Traffic Light Diet^[74] or a variant for dietary prescriptions. Utilizing the USDA's Food Guide Pyramid as its foundation, children are encouraged to increase intake of low-fat nutrient-rich "Green" foods (fruits and vegetables), to consume moderate-calorie "Yellow" foods in moderation (certain grain foods), and minimize if not eliminate high-fat high-sugar "Red" foods (candies). Detailed lists of food alternatives and their corresponding calories are provided to families, who are encouraged to try new and varied Green and Yellow foods. Most behavioral programs initially strive to reduce children's total daily caloric intake while maintaining ade-

quate nutrition for development and growth. A recommended first step is increasing children's awareness of eating habits through self-monitoring, with parental help. With appropriate reductions of total calories and fat intake, children can meet their nutritional needs through increasing the nutrient density of foods eaten, shift toward negative energy balance, and gradually substitute for unhealthy food choices.

The most successful pediatric obesity programs have typically (but not always) included a physical activity component.^[75] There appear to be consistent short-term effects of physical activity interventions on both children's weight status and cardiorespiratory fitness.^[76] Planned aerobic sessions seem to be more beneficial for children's weight loss than lower energy expenditure calisthenics programs;^[75] however, for long-term weight maintenance, data suggest that the best results are achieved through lifestyle approaches that attempt to weave physical activity into every day living (climbing stairs instead of taking the elevator, parking one's car at a distance from a supermarket, walking to the grocery store).^[73,75]

Weight loss can also be achieved by targeting reductions in sedentary activities (TV viewing) rather than targeting increased physical activity per se.^[76-77] These studies show that children who discontinue sedentary activities will naturally redistribute some (although not all) of their time to physical activity which facilitates weight loss. Hence, targeting reduced sedentary activity (TV viewing, video games, computers) has become a vital treatment strategy.

Prevention

The prevention of childhood obesity is one of the most pressing needs in the field. There have been virtually no population-based childhood obesity prevention studies, although several notable school prevention studies have yielded encouraging results. Two of the most influential school-based prevention studies in recent years are those by Robinson^[78] and Gortmaker et al^[79]. Robinson's school-based study compared 192 third- and fourth-grade students (mean age 8.9 years) from two California elementary schools. The intervention consisted of an 18-lesson, 6-month classroom curriculum to reduce television, videotape, and video game use. Compared to the control school, the treatment school demonstrated less of an increase in BMI (intervention vs control changes: 18.38 to 18.67 kg/m² vs 18.10 to 18.81 kg/m²), less of an increase in triceps skin fold measurements (intervention vs control change: 14.55 to 15.47 mm vs 13.97 to 16.46 mm), less of an increase in waist to hip ratio (intervention vs control

change: 0.83 to 0.83 vs 0.82 to 0.84), and less of an increase in waist circumference (intervention vs control change: 60.48 to 63.57 cm vs 59.51 to 64.73 cm). The intervention group also demonstrated statistically significant decreases in reported television viewing (intervention vs control change: 15.35 to 8.80 vs 15.36 to 14.46 hours/week). This study provides strong illustration of the potential role of television viewing reduction in obesity prevention.

Gortmaker et al's^[79] "Planet Health" obesity prevention program also emphasized reduction of television viewing as a central component in their obesity prevention program. The researchers compared five treatment schools against five control schools in a study that included 1295 ethnically diverse children in grades six and seven in Massachusetts, United States. Over two years, the obesity prevalence among girls in the intervention schools declined from 23.6% to 20.3%, while the prevalence among girls in the control schools increased from 21.5% to 23.7%. Moreover, changes in television viewing practices predicted treatment out-

come for girls. Each hour of reduced television watching was independently associated with a 15% reduction in the likelihood of being obese. It should be noted, however, that the control and treatment schools did not differ with respect to obesity incidence (the number of new obesity cases over time).

Recently, James et al^[80] conducted a school-based intervention of 7 to 11-year-old designed to reduce consumption of carbonated beverages. The intervention group reduced their intake of carbonated drinks over 3 days. At 1 year, the percentage of overweight and obese children decreased in the intervention group but increased in the control group.

There is a great need for new and innovative studies for research into prevention of childhood obesity. Based on the results of a recent National Institutes of Health conference on this topic, Kumanyika and Obarzanek^[81] outlined a list of new avenues for research (Table 3). Perusal of the list reveals the amount of work that remains to be done.

Table 3. Directions for future obesity prevention research^[81]

Expand the types of interventions to be tested to prevent obesity and promote weight control and long-term maintenance of weight loss.	<ul style="list-style-type: none"> • Explore approaches such as participatory action research that encourage community input and have an influence on intervention design and content. • Develop and test interventions that target multiple societal levels, individual-based interventions, and their combination, to determine whether public health and individual approaches are interactive. • Develop and test environmental strategies (in homes, organizations, work sites, neighborhoods), including policy changes. • Develop and test family-based interventions and interventions in the primary health care setting, including physician incentives.
Conduct research to improve efficacy of interventions for obesity prevention and long-term weight control programs.	<ul style="list-style-type: none"> • Determine optimal intervention duration, frequency, and mode of delivery (group or individual face-to-face contact, telephone, mail, internet). • Use highly controlled study designs to determine optimal physical activity and dietary prescriptions in adults and in children and adolescents. • Determine the optimal use of meal replacements for weight loss and its long-term maintenance. • Test interactive effects of genetic and/or psychosocial predictors of obesity by stratifying study participants on these factors. • Explore and describe approaches to intervention by internet, alone and in combination with other intervention strategies. Gather information on the "dose" of intervention the internet provides and the ability of the internet to enhance motivation and adherence to interventions.
Conduct observational studies to guide interventions for obesity prevention, weight control, and long-term maintenance of weight loss.	<ul style="list-style-type: none"> • Identify physiological and behavioral characteristics that put individuals at risk for weight gain. • Identify physiological and psychosocial characteristics and other predictors of success in intervention programs. • Identify barriers to enrolling and participating in weight control programs. • Identify approaches and strategies to enhance motivation and adherence to intervention in children and adolescents, women during the prenatal and postpartum period, and other populations.
Use theoretical models to design obesity prevention interventions.	<ul style="list-style-type: none"> • Incorporate theoretical perspectives from other areas of research in the conceptualization and design of obesity prevention studies; for example, family systems theory, and organizational behavior theory. • Conduct studies in which obesity prevention objectives are integrated with other types of health promotion or behavior change objectives. • Develop more systematic concepts of how interventions can be tailored to specific behaviors, populations, and contexts.
Conduct research to improve measurement methodology for obesity prevention research.	<ul style="list-style-type: none"> • Develop valid and affordable energy balance measurements suitable for large-scale trials outside of clinical settings. • Identify psychosocial measures most relevant for obesity research and explore whether these psychosocial measures can be standardized and used in weight control trials. • Determine a health-related and functional definition obesity in children.




Fig. Potential levels for childhood obesity prevention.

Conclusion

A major concern related to childhood obesity is that obese children tend to become obese adults, with all the risks/comorbidities associated (diabetes, cardiovascular diseases, among many others). Efforts to manage and prevent childhood obesity involve education, research and intervention. Research could drive new directions in prevention and develop public policy that might help manage the problem. Additionally, research is needed to test these issues. In conclusion six relevant levels may be involved in the prevention and treatment of pediatric obesity and each of these needs investigation: family, schools, health care, government, industry and media (Fig.). Together these six levels could promote childhood obesity as a high research priority and put it as the first point in the international public agenda.

Funding: This study was funded in part by NIH grant R01-HDO 42169 to MSF.

Ethical approval: Not needed.

Competing interest: None declared.

Contributors: AP wrote the first draft of this paper. All authors contributed to the intellectual content and approved the final version. JK is the guarantor.

References

- 1 International Obesity Task Force data, based on population-weighted estimates from published and unpublished surveys, 1990-2002 using IOTF-recommended cut-offs for overweight and obesity [<http://www.ietf.org>].
- 2 Dietz WH, Robinson TN. Overweight children and adolescents. *N Engl J Med* 2005;352:2100-2109.
- 3 Hedley A, Ogden CL, Johnson CL, Carroll MD, Curtin LR, Flegal KM. Prevalence of overweight and obesity among US children, adolescents, and adults, 1999-2002. *JAMA* 2004; 291:2847-2850.
- 4 Ogden CL, Flegal KM, Carroll MD, Johnson CL. Prevalence and trends in overweight among US children and adolescents. *JAMA* 2002;288:1728-1732.
- 5 Lissau I, Overpeck MD, Ruan WJ, Due P, Holstein BE, Hediger ML. Body mass index and overweight in adolescents in

- 13 European countries, Israel and the United States. *Arch Pediatr Adolesc Med* 2004;158:27-33.
- 6 Wang Y, Monteiro C, Popkin BM. Trends of obesity and underweight in older children and adolescents in the United States, Brazil, China, and Russia. *Am J Clin Nutr* 2002;75: 971-977.
- 7 Kain J, Uauy R, Vio F, Albala C. Trends in overweight and obesity prevalence in Chilean children: comparison of three definitions. *Eur J Clin Nutr* 2002;56:200-204.
- 8 de Onis M, Blossner M. Prevalence and trends of overweight among preschool children in developing countries. *Am J Clin Nutr* 2000;72:1032-1039.
- 9 Gonzales UM, Ramos RI, Martinez AG, Flores HS, Martinez SH. Crecimiento y estado de nutrición. *Atendidos per el Instituto Mexicano del Seguro Social* 2004;153-170.
- 10 Avila Y, Shamah LT, Chavez VA, Galindo GC. Encuesta urbana de alimentacion y nutrition en la zona metropolitana de la Ciudad de Mexico 2002. ENURBAL. Instituto Nacional de Salud Publica, 2003.
- 11 Tremblay MS, Katzmarzyk PT, Williams JD. Temporal trends in overweight and obesity in Canada, 1981-1996. *Int J Obes* 2002;26:538-543.
- 12 Troiano RP, Flegal KM, Kuczmarski RJ, Campbell SM, Johnson CL. Overweight prevalence and trends for children and adolescents. *Arch Pediatr Adolesc Med* 1995;149:1085-1091.
- 13 Flegal KM, Ogden CL, Wei R, Kuczmarski RL, Johnson CL. Prevalence of overweight in US children, comparison of US growth charts from the Center for Disease Control and Prevention with other reference values for body mass index. *Am J Clin Nutr* 2001;73:1086-1093.
- 14 Rolland-Cachera MF, Bellisle F, Deheeger M. Nutritional status and food intake in adolescents living in Western Europe. *Eur J Clin Nutr* 2002;54:S41-46.
- 15 Kromeyer-Hauschild K, Zellner K, Jaeger U, Hoyer H. Prevalence of overweight and obesity among school children in Jena (Germany). *Int J Obes* 1999;23:1143.
- 16 Mamalakis G, Kafatos A, Manios Y, Anagnostopoulou T, Apostolaki I. Obesity indices in a cohort of primary school children in Crete: six year prospective study. *Int J Obes* 2000; 24:765-771.
- 17 Krassas GE, Tzotzas T, Tsamatis C, Konstantinidis T. Prevalence and trends in overweight and obesity among children and adolescents in Thessaloniki, Greece. *J Pediatr Endocrinol Metab* 2001;14:1319-1326.
- 18 Cacciari E, Milani S, Balsamo A, Dammacco F, De Luca F, Chiarelli F, et al. Italian cross-sectional growth charts for height, weight and BMI (6-20y). *Eur J Clin Nutr* 2002;56: 171-180.
- 19 Padez C, Fernandez T, Marques V, Moreira P, Mourao I. Portuguese prevalence study of obesity in childhood: the role of socio-demographic factors. Symposium on Childhood Obesity. Loughborough, UK. December 2003.
- 20 Moreno LA, Sarria A, Fleta J, Rodriguez G, Bueno M. Trends in body mass index and overweight prevalence among children and adolescents in the region of Aragon (Spain) from 1985 to 1995. *Int J Obes* 2000;24:925-931.
- 21 Rios M, Fluiters E, Perez Mendez LF, Garcia-Mayor EG, Garcia-Mayor RV. Prevalence of childhood overweight in North western Spain: a comparative study of two periods with a ten year interval. *Int J Obes* 1999;23:1095-1098.
- 22 Berg IM, Simonsson B, Brantefor B, Ringqvist I. Prevalence of overweight and obesity in children and adolescents in a

- county in Sweden. *Acta Paediatr* 2001;90:671-676.
- 23 Zimmermann MB, Hess SY, Hurrell RF. A national study of the prevalence of overweight and obesity in 6-12-y-old Swiss children: body mass index, body weight perceptions and goals. *Eur J Clin Nutr* 2000;54:568-572.
 - 24 Lobstein TJ, James WPT, Cole TJ. Increasing levels of excess weight among children in England. *Int J Obes* 2003;27:1136-1138.
 - 25 Ibrahim B, Sallam S, Tawila S, Gibaly O, Sahn F. Transitions to adulthood, a national survey of Egyptian adolescents. *The Population Council; New York*;2002;25-30.
 - 26 Magarey AM, Daniels LA, Boulton TJ. Prevalence of overweight and obesity in Australian children and adolescents; reassessment of 1985 and 1995 data against new standard international definition. *Med J Aust* 2001;174:561-564.
 - 27 Ke-You G, Da-Wei F. The magnitude and trends of under- and over-nutrition in Asian countries. *Biomed Environ Sci* 2001;14:53-60.
 - 28 Ramachandran A, Snehalatha C, Vinitha R, Thayil M, Kumar CK, Sheeba L, et al. Prevalence of overweight in urban Indian adolescent school children. *Diabetes Res Clin Pract* 2002;57:185-190.
 - 29 Going SB. Densitometry. In: Roche AF, Heymsfield SB, Lohman TG, eds. *Human Body Composition. Human Kinetics, Champaign, IL*, 1996;3-24.
 - 30 Ma G, Yao M, Liu Y, Lin A, Zou H, Urlando A, et al. Validation of a new pediatric air-displacement plethysmograph for assessing body composition in infants. *Am J Clin Nutr* 2004;79:653-660.
 - 31 Heymsfield SB, Gallagher D, Grams J, Nunez C, Wang ZM, Pietrobelli A. Upper extremity skeletal muscle mass: potential of measurement with single frequency bioimpedance analysis. *Appl Radiat Isot* 1998;49:473-474.
 - 32 Higgins PB, Gower BA, Hunter GR, Goran MI. Defining health-related obesity in prepubertal children. *Obes Res* 2001;9:233-240.
 - 33 Heymsfield SB, Wang ZM, Baumgartner RN, Ross R. Human body composition: advances in models and methods. *Ann Rev Nutr* 1997;17:527-558.
 - 34 Weiss R, Dziura J, Burgert TS, Tamborlane WV, Taksali SE, Yeckel CW, et al. Obesity and the metabolic syndrome in children and adolescents. *N Engl J Med* 2004;350:2362-2374.
 - 35 Goran MI. Measurement issues related to studies of childhood obesity: assessment of body composition, body fat distribution, physical activity, and food intake. *Pediatrics* 1998;101:505-518.
 - 36 Kuczmarski RJ, Ogden CL, Grummer-Starwn LM, Flegal KM, Guo SS, Wei Z, et al. *CDC Growth charts: United States advance data from vital and health statistics, no 314. National Center for Health Statistics; Atlanta*, 2000.
 - 37 Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000;320:1240-1243.
 - 38 Freedman SE, Serdula MK, Srinivasan SR, Berenson GS. The relation of circumferences and skinfolds to levels of lipids and insulin: the Bogalusa Heart Study. *Am J Clin Nutr* 1999;69:308-317.
 - 39 Lohman TG. Assessment of body composition in children. *Pediatr Exerc Sci* 1989;1:19-30.
 - 40 Fernandez J, Pietrobelli A, Redden DT, Allison DB. Waist circumference percentile in nationally representative samples of African-American, European-American, and Mexican-American children and adolescents. *J Pediatr* 2004;145:439-444.
 - 41 Dietz WH. Overweight and precursors of type 2 diabetes mellitus in children and adolescents. *J Pediatr* 2001;138:453-454.
 - 42 Allison DB, Heshka S, Heymsfield SB. Evidence for a major gene with pleiotropic effects for a Cardiovascular Disease Risk Syndrome in children under 14. *Am J Dis Child* 1993;147:1298-1302.
 - 43 Falkner B, Kushner H, Onesti G, Angelakos FT. Cardiovascular characteristic in adolescents who develop essential hypertension. *Hypertension* 1981;3:521-527.
 - 44 Lauer RM, Connor WE, Leaverton PE, Reiter MA, Clarke WR. Coronary heart disease risk factors in school children: the Muscatine Study. *J Pediatr* 1975;86:697-708.
 - 45 Williams DP, Going SB, Lohman TG, Harsha DW, Srinivasan SR, Webber LS, et al. Body fatness and risk for elevated blood pressure, total cholesterol, and serum lipoprotein ratios in children and adolescents. *Am J Public Health* 1992;82:358-362.
 - 46 Gutin B, Basch C, Shea S, Contento I, DeLozier M, Rips J, et al. Blood pressure, fitness, and fatness in 5- and 6-year-old children. *JAMA* 1990;264:1123-1127.
 - 47 Daniels SR, Witt SA, Glascock B, Khoury PR, Kimball TR. Left atrial size in children with hypertension: the influence of obesity, blood pressure, and left ventricular mass. *J Pediatr* 2002;141:186-190.
 - 48 Krebs NF, Jacobson MS. American Academy of Pediatrics Committee on Nutrition. Prevention of pediatric overweight and obesity. *Pediatrics* 2003;112:424-430.
 - 49 Rodriguez MA, Winkleby MA, Ahn D, Sundquist J, Kraemer HC. Identification of population subgroups of children and adolescents with high asthma prevalence: findings from the Third National Health and Nutrition Examination Survey. *Arch Pediatr Adolesc Med* 2002;156:269-275.
 - 50 Baumgartner RN, Siervogel RM, Chumlea WC, Roche AF. Association between plasma lipoprotein cholesterols, adiposity and adipose tissue distribution during adolescence. *J Obes* 1989;13:31-41.
 - 51 Frerichs RR, Webber LS, Srinivasan SR, Berenson GS. Relation of serum lipids and lipoproteins to obesity and sexual maturity in white and black children. *Am J Epidemiol* 1978;108:486-496.
 - 52 Frupp RR, Hodgson IT, Kwiterovich P0, Wernwr JC, Sculsler HG, Whitman V. Aerobic capacity, obesity, and arteriosclerotic risk factors in male adolescents. *Pediatrics* 1985;75:833-848.
 - 53 Smoak CG, Burke GL, Webber LS, Harsha DW, Srinivasan SR, Berenson GS. Relation of obesity to clustering of cardiovascular disease risk factors in children and young adults: the Bogalusa Heart Study. *Am J Epidemiol* 1987;125:364-372.
 - 54 Freedman DS, Srinivasan SR, Harsha DW, Webber LS, Berenson GS. Relation of body fat patterning to lipid and lipoprotein concentrations in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr* 1989;50:930-939.
 - 55 Bandini LG, Schoeller DA. Total body fat distribution and glucose tolerance in adolescents. *Am J Clin Nutr* 1986;43:696.
 - 56 Freedman DS, Srinivasan SR, Burke GL, Shear CL, Smoak CG, Harsha DW, et al. Relation of body fat distribution to hyperinsulinemia in children and adolescents: the Bogalusa Heart Study. *Am J Clin Nutr* 1987;46:403-410.
 - 57 Jacot JP, Zuppinger KA, Joss EE, Donatin A. Evidence for two types of juvenile obesity on the basis of body composition and insulin release following small doses of glucose. *Klinische Wochenschrift* 1973;51:1109-1114.

- 58 Jiang X, Srinivasan SR, Bao W, Berenson GS. Association of fasting insulin with blood pressure in young individuals. *Arch Intern Med* 1993;153:323-328.
- 59 Voors AW, Harsha DW, Webber LS, Radhakrishnamurthy B, Srinivasan SR, Berenson GS. Clustering of anthropometric parameters, glucose tolerance, and serum lipids in children with high and low beta-and pre-beta-lipoproteins. *Bogalusa Heart study. Arteriosclerosis* 1982;2:346-355.
- 60 Modan M, Halkin H, Almog S, Lusky A, Eshkol A, Shefi M, et al. Hyperinsulinemia: a link between hypertension, obesity and glucose intolerance. *J Clin Invest* 1985;75:809-817.
- 61 Rocchini AP, Katch V, Kveselis D, Moorehead C, Martin M, Lampman R, et al. Insulin and renal retention in obese adolescents. *Hypertension* 1989;14:367-374.
- 62 Sinaiko AR, Donahue RP, Jacobs DR Jr, Prineas RJ. Relation of weight and rate of increase in weight during childhood and adolescence to body size, blood pressure, fasting insulin, and lipids in young adults. *The Minneapolis Children's Blood Pressure Study. Circulation* 1999;11:1471-1476.
- 63 Goran MI. Metabolic precursors and effects of obesity in children: a decade of progress, 1990-1999. *Am J Clin Nutr* 2001;73:158-171.
- 64 Sinaiko AR, Jacobs DR Jr, Steinberger J, Moran A, Luepker R, Rocchini AP, et al. Insulin resistance syndrome in childhood: associations of the euglycemic insulin clamp and fasting insulin with fatness and other risk factors. *J Pediatr* 2001;139:700-707.
- 65 Pinhas-Hamiel O, Dolan LM, Daniels SR. Increased incidence of non-insulin-dependent diabetes mellitus among adolescents. *J Pediatr* 1996;128:608-615.
- 66 Coates TJ, Jeffery RW, Slinkard LA, Killen JD, Danaher BG. Frequency of contact and monetary reward in weight loss, lipid change, and blood pressure reduction with adolescents. *Behav Ther* 1982;13:175-185.
- 67 Epstein LH, Wing RR, Steranchak L, Dickson B, Michelson J. Comparison of family-based behavior modification and nutrition education for childhood obesity. *J Pediatr Psychol* 1980;5:25-36.
- 68 Johnson WG, Hinkel LK, Carr RE, Anderson DA, Lemmon CR, Engler LB, et al. Dietary and exercise interventions for juvenile obesity: long-term effects of behavioral and public health models. *Obes Res* 1997;5:257-261.
- 69 Epstein LH, Wing RR, Penner BC, Kress MJ. Effect of diet and controlled exercise on weight loss in obese children. *J Pediatr* 1985;107:358-361.
- 70 Rocchini AP, Katch V, Anderson J, Hinderliter J, Becque D, Martin M, et al. Blood pressure in obese adolescents: effect of weight loss. *Pediatrics* 1988;82:16-23.
- 71 Epstein LH, Valoski A, Wing RR, McCurley J. Ten-year outcomes of behavioral family-based treatment for childhood obesity. *Health Psychol* 1994;13:373-383.
- 72 Blomquist B, Borjeson M, Larsson Y, Persson B, Sterky G. The effect of physical activity on the body measurements and work capacity of overweight boys. *Acta Paediatr Scand* 1965;54:566-572.
- 73 Epstein LH, Wing RR, Koeske R, Ossip D, Beck S. A comparison of lifestyle change and programmed aerobic exercise on weight and fitness changes in obese children. *Behav Ther* 1982;13:651-665.
- 74 Epstein LH, Squires S. *The stoplight diet for children: an eight-week program for parents and children*. Boston: Little, Brown and Company, 1988.
- 75 Epstein LH, Wing RR, Koeske R, Valoski A. A comparison of lifestyle exercise, aerobic exercise, and calisthenics on weight loss in obese children. *Behav Ther* 1985;16:345-356.
- 76 Epstein LH, Valoski AM, Vara LS, McCurley J, Wisniewski L, Kalarchian MA, et al. Effects of decreasing sedentary behavior and increasing activity on weight change in obese children. *Health Psychol* 1995;14:109-115.
- 77 Epstein LH, Paluch RA, Gordy CC, Dorn J. Decreasing sedentary behaviors in treating pediatric obesity. *Arch Pediatr Adolesc Med* 2000;154:220-226.
- 78 Robinson TN. Reducing children's television viewing to prevent obesity: a randomized controlled trial. *JAMA* 1999;282:1561-1567.
- 79 Gortmaker S, Peterson K, Wiecha J, Sobol AM, Dixit S, Fox MK, et al. Reducing obesity via a school-based interdisciplinary intervention among youth: Planet Health. *Arch Pediatr Adolesc Med* 1999;153:409-418.
- 80 James J, Thomas P, Cavan D, Kerr D. Preventing childhood obesity by reducing consumption of carbonated drinks: cluster randomized controlled trial. *BMJ* 2004;328:1237.
- 81 Kumanyika SK, Obarzanek E. Pathways to obesity prevention: report of a national institutes of health workshop. *Obes Res* 2003;11:1263-1274.

Received June 19, 2005

Accepted after revision June 22, 2005